arXiv:hep-th/9410183v1 25 Oct 1994

hep-th/9410183
MPI-PhT/94-66
ZU-TH 32/94
October 1994

# BLACK HOLES AND SPHALERONS IN LOW ENERGY EFFECTIVE STRING THEORY [*]

GEORGE LAVRELASHVILI [†] [‡]

*Institut für Theoretische Physik, Universität Zürich*
*Winterthurerstrasse 190, CH-8057 Zürich, Switzerland* [§]

and

*Max-Planck-Institut für Physik, Werner-Heisenberg-Institut*
*Föhringer Ring 6, D-80805 Munich, Germany*

## ABSTRACT

We discuss properties of globally regular and black hole solutions of the Einstein-Yang-Mills-dilaton theory.

---

[*] Talk given at the 7th Marcel Grossmann Meeting on General Relativity, Stanford University, July 24-30, 1994. To be published in the Proceedings.

[†] Electronic address: lavrela@physik.unizh.ch

[‡] On leave of absence from Tbilisi Mathematical Institute, 380093 Tbilisi, Georgia

[§] Address after October 1, 1994.

The existence of a dilaton - a neutral, scalar field with exponential coupling to matter is one of the universal predictions following from various types of string theories and from Kaluza-Klein theories (see e.g. [1]). The dilaton field has an influence on the cosmological evolution of the Universe [2] and it might modify particle-like solutions of corresponding theories.

In the present report we are interested in the second aspect of the problem, namely we will study static spherically symmetric solutions of the Einstein-Yang-Mills-dilaton (EYMD) theory. We will consider the EYMD theory defined by the action

$$S = \frac{1}{4\pi}\int\Big(-\frac{1}{4G}R + \frac{1}{2}(\partial\varphi)^2 - \frac{\mathrm{e}^{2\kappa\varphi}}{4g^2}F^2\Big)\sqrt{-g}d^4x \ , \tag{1}$$

where $G$ is Newton's constant, $g$ is the gauge coupling constant and $\kappa$ denotes the dilatonic coupling constant. The scaling properties of the action Eq.(1) allow us to put $G = g = 1$ without restrictions in what follows. The only remaining free parameter in the action is the dimensionless ratio $\gamma = \kappa/g\sqrt{G}$.

Note that in the limit $\gamma \to 0$ the model described by Eq. (1) reduces to the Einstein-Yang-Mills theory [3, 4]. In the limit $\gamma \to \infty$ one gets the Yang-Mills-dilaton theory [5] in flat space.

To study static, spherically symmetric solutions of EYMD theory a convenient parametrization for the metric turns out to be

$$ds^2 = A^2(r)\mu(r)dt^2 - \frac{dr^2}{\mu(r)} - r^2 d\Omega^2 \ , \tag{2}$$

where $d\Omega^2 = d\theta^2 + sin^2(\theta)d\varphi^2$ is the line element of the unit sphere.

For the $SU(2)$ Yang-Mills potential we make the usual ('magnetic') spherically symmetric ansatz

$$W_0^a = 0, \quad W_i^a = \epsilon_{aik}\frac{x^k}{r^2}(W(r) - 1) \ . \tag{3}$$

Substituting this ansatz into the action we obtain the reduced action

$$S_{\mathrm{red}} = -\int A\Big[\frac{1}{2}(\mu + r\mu' - 1) + \frac{r^2}{2}\mu\varphi'^2 + \mathrm{e}^{2\gamma\varphi}\Big(\mu W'^2 + \frac{(1 - W^2)^2}{2r^2}\Big)\Big]\, dr \ , \tag{4}$$

where a prime denotes $\frac{d}{dr}$.

The resulting field equations are

$$
\begin{aligned}
(A\mathrm{e}^{2\gamma\varphi}\mu W')' &= A\mathrm{e}^{2\gamma\varphi}\frac{W(W^2-1)}{r^2} \ , \\
(A\mu r^2\varphi')' &= 2\gamma A\mathrm{e}^{2\gamma\varphi}\Big(\mu W'^2 + \frac{(1-W^2)^2}{2r^2}\Big) \ , \\
\mu' &= \frac{1}{r}\bigg(1-\mu-r^2\mu\varphi'^2 - 2\mathrm{e}^{2\gamma\varphi}\Big(\mu W'^2 + \frac{(1-W^2)^2}{2r^2}\Big)\bigg) \ , \\
A^{-1}A' &= \frac{2\mathrm{e}^{2\gamma\varphi}W'^2}{r} + r\varphi'^2 \ . \qquad (5)
\end{aligned}
$$

The field equations Eq. (5) have singular points at $r=0$ and $r=\infty$ as well as at points where $\mu(r)$ vanishes.

Inserting a power series expansion into Eq. (5) one finds a 2−parameter family of regular solutions in the vicinity of $r=0$

$$
\begin{aligned}
W(r) &= 1-br^2+O(r^4) \ , \qquad \mu(r) = 1-4b^2\mathrm{e}^{2\gamma\varphi_0}r^2+O(r^4) \ , \\
\varphi(r) &= \varphi_0 + 2\gamma\mathrm{e}^{2\gamma\varphi_0}b^2r^2+O(r^4) \ , \quad A(r) = 1+4b^2\mathrm{e}^{2\gamma\varphi_0}r^2+O(r^4) \ , (6)
\end{aligned}
$$

where $b$ and $\varphi_0$ are arbitrary parameters.

Similarly at $r=\infty$ one finds

$$
\begin{aligned}
W(r) &= \pm(1-\frac{c}{r}+O(\frac{1}{r^2})) \ , \quad \mu(r) = 1-\frac{2M}{r}+O(\frac{1}{r^2}) \ , \\
\varphi(r) &= \varphi_\infty - \frac{d}{r}+O(\frac{1}{r^2}) \ , \quad A(r) = A_\infty(1-\frac{d^2}{2r^2}+O(\frac{1}{r^4})) \ , \qquad (7)
\end{aligned}
$$

where again $c, d, M, \varphi_\infty$ and $A_\infty$ are arbitrary parameters.

In the vicinity of the singular point $r_h$ where $\mu(r)$ vanishes one finds a 2−parameter family of solutions which stay regular at this point. Under an appropriate choice of the parameters the surface $r=r_h$ corresponds to a regular event horizon.

Globally regular asymptotically flat (respectively back hole) solutions have to interpolate between the described asymptotic behavior at $r=0$ (respectively $r=r_h$) and $r=\infty$.

The analysis [6, 7] for $0<\gamma<\infty$ yields what could be expected from the extreme cases $\gamma=0$ and $\gamma=\infty$. It was found [6] that for any value of the dilaton coupling constant $\gamma$ the EYMD system has: a discrete family of globally regular (sphaleron type) solutions of finite mass and a discrete family of (non-Abelian) black hole solutions. The solutions in each family are labeled by the number $n$ of zeros of the gauge field potential $W(r)$.

The solutions are found to be unstable [6].

It is interesting that the black holes have a non-vanishing Yang-Mills field outside the horizon. One can interpret the solutions with the horizon as "black holes inside sphalerons".

The mass of the solutions in the case of the Einstein-Yang-Mills theory is of the order of one in natural units $M_{EYM} = 1/g\sqrt{G}$. In the EYMD theory the mass of the solutions decreases with increasing dilatonic coupling constant $\gamma$ and for large $\gamma$ goes to zero like $M_{EYMD} \sim \frac{M_{EYM}}{\gamma^2}$.

Globally regular solutions with odd $n$ are kinds of sphalerons. One finds fermion zero modes in the background of these solutions [8, 9, 10]. One can assign a topological number to these solutions. It is half integer.

There are some interesting regularities in the EYMD theory. It turns out that the $tt$ component of the metric is related to the dilaton $A^2\mu = e^{2\gamma\varphi}$ and as a consequence of this relation the dilaton charge (parameter $d$ in Eq.(7)) is equal to the mass of the solution: $d = M$.

A very special situation occurs for the value of the dilaton coupling constant $\gamma = 1$, which corresponds to the model obtained from heterotic string theory. It was found [6] that for the $n = 1$ solution the parameter $b$ is a rational number, $b = \frac{1}{6}$. Another regularity found numerically is that an asymtotic coefficient $c$ in Eq.(7) is related to the mass of the solution, $c = 2M$. We think these are arguments indicating that the lowest lying ($n = 1$) regular solution may be obtained in closed form. similarly to the "stringy instanton" and the "stringy monopole"[11].

Due to the high mass of the solutions the only situation where they could play a role is in the Early Universe, but at the moment there seems to be no natural physical scenario where we could make use of these solutions.

# Acknowledgements

I am grateful to Dieter Maison for a fruitful collaboration which led to the results reported in this paper.

# References

[1] M.B. Green, J.H. Schwarz and E. Witten, *Superstring Theory* (Cambridge U.P., Cambridge, 1987).

[2] See e.g.,
O. Bertolami, Talk presented at the MG7;
M.C. Bento and O. Bertolami, *Phys. Lett.* **B336** (1994) 6.

[3] R. Bartnik and J. McKinnon, *Phys. Rev. Lett.* **61** (1988) 141.

[4] M.S. Volkov and D.V. Gal'tsov, *JETP Lett.* **50** (1990) 346;
H.P. Künzle and A.K.M. Masood-ul-Alam, *J. Math. Phys.* **31** (1990) 928;
P. Bizon, *Phys. Rev. Lett.* **61** (1990) 2844.

[5] G. Lavrelashvili and D. Maison, *Phys. Lett.* **B295** (1992) 67;
P. Bizon, *Phys. Rev.* **D47** (1993) 1656.

[6] G. Lavrelashvili and D. Maison, *Nucl. Phys.* **B410** (1993) 407;
G. Lavrelashvili and D. Maison, *Dilatonic sphalerons and non-abelian black holes,* Preprint MPI-Ph/93-55, hep-th/9307159.

[7] E.E. Donets and D.V. Gal'tsov, *Phys. Lett.* **B302** (1993) 411;
P. Bizon, *Acta Physica Polonica* **B24** (1993) 1209;
T. Torii and Kei-ichi Maeda, *Phys. Rev.* **D48** (1993) 1643.

[8] G.W. Gibbons and A. Steif, *Phys. Lett.* **B314** (1993) 13.

[9] M.S. Volkov, *Phys. Lett.* **B334** (1994) 40.

[10] G. Lavrelashvili, *Fermions in the background of dilatonic sphalerons*, hep-th/9410178.

[11] C.G. Callan, J.A. Harvey and A. Strominger, *Nucl. Phys.* **B359** (1991) 611;
J.A. Harvey and J.Liu, *Phys. Lett.* **B268** (1991) 40.